

12916 P06
CAB/mej
8/3/2001

Express Mail Label: EL449700769US

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

APPLICATION FOR LETTERS PATENT
(UTILITY PATENT)

APPLICANT:

R. Preston Mason, a resident of Beverly, MA,
a U.S. citizen

POST OFFICE ADDRESS:

c/o Elucida Research LLC
100 Cummings Center Suite 135L
Beverly, Mass. 01915

INVENTION TITLE:

SYNERGISTIC EFFECT OF
ANLODIPINE AND ATORVASTATIN ON
CHOLESTEROL CRYSTAL FORMATION
INHIBITION AND AORTIC
ENDOTHELIAL CELL NITRIC OXIDE
RELEASE

ATTORNEYS:

Jerry Cohen (Reg. No. 20,522)
Harvey Kaye* (Reg. No. 18,978)
Stephen Y. Chow (Reg. No. 31,338)
Christine M. Kuta, (Reg. No. 38,001)
Jacob N. Erlich (Reg. No. 24,338)
Dr. Chester A. Bisbee, (Reg. No. 44,538)
Peter J. Borghetti (Reg. No. 42,345)
Katherine E. Noll (Reg. No. 48811)
Perkins, Smith & Cohen
One Beacon Street
Boston, Massachusetts 02108
(617) 854-4000

* Mr. Kaye is available at 301-948-5535

TO: Honorable Assistant Commissioner of Patents
Washington, D.C. 20231

Sir:

Your applicant, named above hereby petitions grant of a utility patent to him or any assignee of record, at the time of issuance, for an invention more particularly described in the following specification and claims, with the accompanying drawings, verified by the accompanying Declaration and entitled:

SYNERGISTIC EFFECT OF AMLODIPINE AND ATORVASTATIN ON CHOLESTEROL CRYSTAL FORMATION INHIBITION AND AORTIC ENDOTHELIAL CELL NITRIC OXIDE RELEASE

5

Cross-Reference to Related Applications

This application claims priority from U.S. Provisional Patent Application No. 60/223,214
entitled "Synergistic Effect of Amlodipine and Atorvastatin on the Stimulation of Nitric Oxide
10 Release from Rabbit Aorta Endothelial Cells" that was filed on August 4, 2000.

Field of the Invention

15 This invention relates to the effect of amlodipine and atorvastatin, alone and in
combination, on cholesterol crystal formation and the release of nitric oxide (NO) from
endothelial cells.

Background of the Invention

Coronary artery disease (CAD) is the leading cause of mortality in the developed world,
and is associated with substantial morbidity as well. Typically, the patient with CAD has several
20 concomitant conditions, including hypertension, diabetes, and dyslipidemia, increasing overall
risk for poor outcomes and complicating treatment. A therapeutic goal for the treatment of CAD
is the development of drugs that can simultaneously target multiple underlying disease processes
that contribute to atherosclerosis, thereby altering the course of the disease. Therefore, CAD
therapy may have increased positive outcomes if the use of an antihypertensive agent and HMG-
25 CoA reductase inhibitor was combined in a single delivery system.

Free cholesterol is an important structural component of the cell plasma membrane that modulates packing of phospholipid molecules, thus regulating lipid bilayer dynamics and structure. The cholesterol molecule is oriented in the membrane such that the long-axis lies parallel to the phospholipid acyl chains, increasing order in the upper acyl chain region of the membrane while decreasing packing constraints at the terminal methyl groups. During atherogenesis, however, increasing levels of cellular cholesterol lead to its abnormal deposition in the vessel wall and the formation of cholesterol crystals.

In animal models of atherosclerosis, it has been demonstrated that the cholesterol content of membranes associated with vascular smooth muscle and macrophage foam cells becomes elevated, resulting in the formation of discrete domains. These highly organized cholesterol structures, characterized by a unit cell periodicity of 34.0 Å, appear to serve as nucleating sites for the formation of extracellular crystals. These domains have been previously described in model membrane systems. A recent study from our laboratory showed that cultured mouse peritoneal macrophage foam cells produced free cholesterol crystals that extend from intracellular membrane sites with various morphologies that include plates, needles and helices. With the use of x-ray diffraction approaches, the early stages of crystal formation could be identified in isolated membranes from these cells. Preventing crystal formation is an important goal as cholesterol in this state is practically inert and does not respond well to pharmacologic interventions that promote lesion regression.

In addition, the normal production of NO by the endothelium is critical for maintaining vascular function. During atherosclerosis, however, endothelial dysfunction effects a significant

reduction in NO production, resulting in: 1) increased monocyte and LDL infiltration, 2) loss of smooth muscle cell function and abnormal proliferation, 3) increased oxidative stress, and 4) increased platelet aggregation. Pharmacologic interventions that restore endothelial function and NO metabolism have demonstrated benefit in the treatment of various cardiovascular disorders, including coronary artery disease.

However, no pharmaceutical composition currently exists that treats both hypertension and hyperlipidemia. Such a pharmaceutical composition would have several benefits. For example, the multiple risk factors for arterial and related heart disease that are often present in an individual patient could be targeted simultaneously. Additionally, the ease of taking one combined dosage could significantly enhance patient compliance with therapeutic regimens.

Therefore, it is an object of this invention to provide a combination therapy that will treat the multiple pathological processes involved in arterial and related heart disease. These include, but are not limited to, hypertension and hyperlipidemia. It is also an object of this invention to develop useful and convenient dosage levels and forms of such a combination therapeutic. Preferably, this pharmaceutical composition would have synergistic effects on these hallmarks of arterial and related heart disease, such that the individual effects of the components of this composition would be enhanced by their combination.

20

Thus, this invention encompasses a therapeutic goal for the treatment of CAD that entails the development of drugs that can simultaneously target multiple underlying disease processes that contribute to atherosclerosis, thereby altering the course of the disease. Therefore, using this

invention, CAD therapy may have increased positive outcomes if the use of an antihypertensive agent and HMG-CoA reductase inhibitor was combined in a single delivery system.

Summary of the Invention

5 Lipophilic agents that influence the organization of lipids may interfere with the early formation of cholesterol domains within the membrane, thereby disrupting crystal formation. The charged dihydropyridine amlodipine (AML) was used for this study as it has strong affinity for phospholipid constituents of atherosclerotic-like membranes, as compared to other calcium channel blockers (CCBs). This property of the drug is attributed to a distinct membrane location
10 that facilitates strong electrostatic binding to the phospholipid phosphate groups, as indicated by x-ray diffraction, differential scanning calorimetry and nuclear magnetic resonance analyses. These distinct membrane interactions may underlie the ability of AML to reverse cholesterol-induced increases in the width of atherosclerotic membranes obtained from vascular smooth muscle cells and to interfere with oxidative modification of lipid constituents.

15 The various membrane biophysical effects of AML may contribute to its observed benefit in coronary artery disease, as demonstrated in the Prospective Randomized Evaluation of the Vascular Effects of Norvasc Trial (PREVENT). The results of PREVENT showed a reduction in cardiovascular morbidity associated with AML therapy, including a 42% reduction in the need
20 for revascularization and 33% reduction in documented angina, as compared to placebo. AML therapy was also associated with significant slowing in the progression of carotid atherosclerosis, as measured with B-mode ultrasonographic assessments. As such clinical benefit has not been

observed with other CCB analogs in similarly designed trials, it has been proposed that AML may have distinct antiatherosclerotic activity.

Hypolipidemic therapy has also been demonstrated to be very useful in reducing morbidity and mortality associated with CAD. Among HMG-CoA reductase inhibitors, atorvastatin calcium (AT) has been shown to be very effective as hypolipdemic therapy. In a recent clinical study involving patients with stable and advanced CAD, aggressive lipid-lowering therapy with AT significantly delayed the time to a first ischemic event and reduced, by 36%, the overall incidence of cardiovascular events, as compared to angioplasty with usual medical care.

Small angle x-ray diffraction approaches were utilized to examine the separate and combined effects of AML and AT on the formation of cholesterol monohydrate structures (*d*-space of 34.0 Å) within the membrane bilayer. This question was investigated in membranes containing cholesterol at levels that reproduce atherosclerotic-like conditions. The results of this study demonstrated that these compounds interfered with the aggregation of cholesterol into separate intrabilayer domains in an unexpected and highly synergistic fashion. This observed effect could not be reproduced by the combination of these compounds with other related drugs. By disrupting the development of cholesterol crystals, this drug combination may facilitate the removal of excess sterol from the vessel wall by plasma high-density lipoproteins (HDL).

In addition, NO production from endothelial cells in rabbit aorta was systematically measured in the absence and presence of amlodipine, atorvastatin, and the combination of the two compounds. The results of these experiments demonstrated a dramatic synergistic effect of

these compounds on NO stimulation. These findings indicate that these agents may effectively stabilize, in a synergistic fashion, NO in the cell by interfering with oxidative destruction.

Other objects, features, and advantages of the present invention will be apparent from the following Detailed Description of the Preferred Embodiments taken in conjunction with the accompanying drawings in which:

Brief Description of the Drawings

Figure 1 shows the X-ray diffraction pattern and corresponding molecular model for cholesterol-enriched membrane bilayer. Diffraction peaks corresponding to sterol-rich and -poor domains can be clearly distinguished at 87% relative humidity at 20°C. The peaks labeled 1' and 2' correspond to the sterol-rich domain ($d = 34.0 \text{ \AA}$) while the surrounding sterol-poor area of the membrane had a d -space value of 60.7 \AA , corresponding to peaks labeled 1, 2 and 4. The corresponding molecular model demonstrates cholesterol bilayer domain with a dimension of 34.0 \AA (each individual cholesterol monohydrate molecule is 17.0 \AA) that is highlighted by the shaded region of the figure.

Figure 2 shows the differential effects of temperature (Figure 2A) and relative humidity (Figure 2B) on the molecular dimensions of cholesterol monohydrate domains versus surrounding sterol-poor membrane regions for samples containing verapamil. The membrane width, as measured in \AA units by x-ray diffraction analysis, represents the distance from the center of one membrane to the next, including surface hydration. In Figure 2A, the effect of temperature on membrane width was evaluated at a constant 93% relative humidity while in

Figure 2B the effect of relative humidity was measured at a constant temperature of 20°C. These data demonstrate that the structure of the cholesterol monohydrate crystalline domains (34.0 Å) are unaffected by changes in temperature or humidity, as compared to the surrounding sterol-poor region of the membrane.

5

Figure 3 shows the X-ray diffraction pattern from oriented membrane lipid bilayers containing elevated levels of cholesterol (1.1:1 and 1.2:1 cholesterol to phospholipid mole ratios) prepared in the absence or presence of the AML/AT combination at 5°C. At a 1.1:1 cholesterol to phospholipid mole ratio, peaks labeled 1, 2 and 4 correspond to *d*-space values of 54.2 Å and 53.0 Å, respectively, for the control and drug-containing samples. At a 1.2:1 cholesterol to phospholipid mole ratio, peaks labeled 1 and 2 corresponded to *d*-space values of 55.5 Å and 53.5 Å, respectively, for the control and drug-containing samples. This figure demonstrates that at a low concentration (30 nM), the combination of AML and AT completely blocked the aggregation of cholesterol into discrete cholesterol domains.

10

15

20

Figure 4 shows the X-ray diffraction patterns from oriented membrane lipid bilayers containing elevated levels of cholesterol (1.2:1 cholesterol to phospholipid mole ratio) prepared in the absence or presence of AML alone, AT alone, AML/AT combination, AT/nifedipine combination, and AML/lovastatin combination at 5°C. The peaks labeled 1, 2 and 4 correspond to the sterol-poor region of the membrane while peaks labeled 1' and 2' correspond to the structure of cholesterol monohydrate domains within the membrane (34.0 Å). The dimensions of the surrounding sterol-poor regions were as follows: control (55.5 Å), AML alone (57.8 Å), AT alone (56.8 Å), AML/AT (53.5 Å), AT/nifedipine (56.5 Å) and AML/lovastatin (54.4 Å).

These experiments demonstrated that the ability of the AML/AT combination to interfere with membrane cholesterol domain formation could not be reproduced by the drugs separately or other CCB/statin combinations.

5 Figure 5 shows the X-ray diffraction patterns from oriented membrane lipid bilayers containing elevated levels of cholesterol (1.1:1 cholesterol to phospholipid mole ratio) prepared in the absence or presence of AML alone, AT alone, and AML/AT combination at 5°C. The peaks labeled 1, 2 and 4 correspond to the sterol-poor region of the membrane while peaks labeled 1' and 2' correspond to the structure of cholesterol monohydrate domains within the membrane (34.0 Å). The dimensions of the surrounding sterol-poor regions were as follows: control (52.4 Å), AML alone (54.4 Å), AT alone (55.8 Å), and AML/AT (53.9 Å). These experiments demonstrated that the AML/AT combination was able to interfere with membrane cholesterol domain formation in a manner that could not be reproduced by the drugs separately.

15 Figure 6 shows the effect of amlodipine, atorvastatin (Compound T), and a mixture of the two compounds on NO release from a single rabbit aorta endothelial cell.

20 Figure 7 shows the dose response curves for NO release stimulated by amlodipine, atorvastatin (Compound T), and a mixture of amlodipine with varying concentrations of atorvastatin (Compound T).

Detailed Description of the Preferred Embodiments

Experimental Methods:

Preparation of reconstituted membrane samples. Porcine cardiac phospholipid dissolved in HPLC-grade chloroform (10.0 mg/ml) was obtained from Avanti Polar Lipids Inc. (Alabaster, AL) and stored at -80°C. The fatty acid composition of the phosphatidylcholine lipids was determined by gas-liquid chromatographic analysis. The overall ratio of saturated to unsaturated fatty acids was 0.8:1, with the primary constituents being 18:2 linoleic acid (30%), 16:0 palmitic acid (22%), 18:1 oleic acid (13%), and 20:4 arachidonic acid (11%). Cholesterol powder was also purchased from Avanti Polar Lipids Inc. Amlodipine besylate (AML) was obtained from Pfizer Central Research (Groton, CT) while atorvastatin calcium (AT) was provided by Parke Davis (Ann Arbor, MI).

The effects of the drugs on membrane cholesterol organization and structure were assessed in well-defined lipid vesicles containing equimolar levels of cholesterol and phospholipid. This reconstituted membrane system was used for the following reasons: 1) this system reproduces changes in membrane structure observed in cholesterol-enriched, atherosclerotic macrophage and smooth muscle cell membranes, 2) the membrane preparation does not contain calcium channels, and 3) these samples can be prepared in a highly reproducible fashion. Lipid vesicles were formed from phospholipid and cholesterol dissolved in chloroform at a fixed molar ratio and added to individual glass 13 x 100-mm test tubes. The chloroform solvent was removed by shell-drying under a steady stream of N₂ gas. Residual solvent was removed under vacuum while the samples were shielded from light. Membrane vesicles were produced for diffraction analysis by rapidly mixing the dried lipids at room temperature

following addition of buffered saline (0.5 mmol/L HEPES and 154.0 mmol/L NaCl, pH, 7.2). The final phospholipid concentration was 5.0 mg/ml. Membrane samples were oriented for diffraction analysis by centrifugation and then placed in hermetically sealed canisters that controlled temperature and relative humidity, as previously described.

5

Small angle x-ray diffraction analysis. Small-angle x-ray diffraction approaches were used to directly examine the effects of the various drugs on the organization of cholesterol in the membrane. X-ray diffraction experiments were conducted by aligning the samples at grazing incidence with respect to a collimated, nickel-filtered monochromatic x-ray source ($\text{CuK}_\alpha = 1.54 \text{ \AA}$) produced by a high-brilliance rotating anode microfocus generator (Rigaku Rotaflex RU-200, Danvers, MA). The diffraction data were collected on a one-dimensional, position-sensitive electronic detector (Innovative Technologies, Newburyport, MA) placed at a distance of 150 mm from the sample. In addition to direct calibration of the detector system, cholesterol monohydrate crystals were used to verify the calibration, as previously described. The unit cell periodicity, or d -space, of the membrane lipid bilayer is the measured distance from the center between one bilayer to the next, including surface hydration, and calculated from Bragg's Law.

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
217

electrode)]. The three electrodes were connected to a potentiostat/galvanostat PAR273. Data were acquired with the use of an IBM computer with custom software. The current proportional to NO concentration was measured by porphyrinic sensor, which operated in amperometric mode at constant potential of 0.63 V vs. SCE.

5

The release of NO was initiated by the injection of potential agonists of endothelial NO synthase (eNOS) using a temtoinjector placed in the controlled distance from the endothelial cell. Two different agonists were tested: amlodipine and atorvastatin. The different concentrations of these two compounds applied simultaneously were also tested.

10

15

Atherosclerotic-like membranes have distinct crystalline-like sterol domains: Membrane sterol-rich domains may represent an important nucleating site for free cholesterol crystal formation, an important feature of the unstable plaque. The separate and combined effects of AML and AT on cholesterol monohydrate formation in membranes reconstituted from native phospholipids isolated from cardiac tissue was evaluated. Phospholipid composed of heterogeneous acyl chains was used for these analyses. This membrane system reproducibly formed discrete sterol-rich domains at levels previously observed in atherosclerosis studies under similar experimental conditions.

20

X-ray diffraction analysis of oriented, cholesterol-enriched membranes produced strong, reproducible diffraction orders that correspond to structurally distinct sterol-rich and -poor membrane regions. The *d*-space measurement refers to the average distance from the center of one membrane bilayer to the next, including surface hydration. The *d*-space of the sterol-rich

region was 34.0 Å, indicative of a cholesterol bilayer structure as a single cholesterol monohydrate molecule has a long axis of 17 Å (Figure 1). The surrounding sterol-poor regions, meanwhile, had an average width of 65.9 Å at 20°C and 93% relative humidity. The much larger width (>90 %) of the sterol-poor domains is attributed to the abundance of phospholipid in the surrounding membrane region. The cholesterol domains were invariably present over a wide range of temperatures (5-37°C) and relative humidity levels (74-93%), consistent with previous x-ray diffraction analyses on atherosclerotic-like membrane samples.

In Figure 1, diffraction peaks corresponding to the sterol-rich and -poor domains can be clearly distinguished at 20°C. The peaks labeled 1' and 2' correspond to the sterol-rich domain ($d = 34.0$ Å) while the surrounding sterol-poor area of the membrane had a d -space value of 60.7 Å, corresponding to peaks labeled 1, 2 and 4. The peaks that describe the cholesterol monohydrate phase are very sharp, as expected for a crystalline-like structure. In every sample that was evaluated, it was observed that the dimensions of the sterol-poor region of the membrane was modulated by temperature and relative humidity due to its heterogeneous chemical composition and the dynamic mobility of the phospholipid-cholesterol binary mixture. At 93% relative humidity, for example, the d -space of the sterol-poor region decreased by 5.5 Å (9%) as sample temperature was increased from 15°C (64 Å) to 40°C (58.5 Å), consistent with increased *trans-gauche* isomerizations (Figure 2). Over this same temperature range, however, the cholesterol monohydrate phase remained unchanged at 34.0 Å, as expected for a crystalline-like structure. In addition, the highly reproducible 34.0 Å structure was unaffected by large changes in relative humidity (52 to 93%) at 20°C while the sterol-poor region changed by 19% or 10 Å (52 to 62 Å) over this same range.

Synergistic inhibition of sterol domain formation with amlodipine and atorvastatin: The

addition of both AML and AT to cholesterol-enriched membrane samples prevented sterol domain formation in a synergistic fashion. At an aqueous buffer concentration of 30 nM, the combination of AML and AT completely blocked the formation of cholesterol domains in membrane samples containing cholesterol and phospholipid at 1.1:1 and 1.2:1 cholesterol:phospholipid mole ratios. In the presence of the two drugs, only peaks corresponding to the phospholipid bilayer could be observed under a variety of experimental conditions, as compared to control (Figure 3). At a 1.1:1 mole ratio, the d -space values for the control and drug combination-containing samples were 54.2 and 53.9 Å, respectively, at 74% relative humidity and 5°C. At a 1.2:1 mole ratio, the d -space values for the control and drug combination-containing samples were 55.5 and 53.5 Å, respectively, at 74% relative humidity and 5°C.

When AML or AT were added separately to the membrane samples, cholesterol domains could be clearly detected under identical conditions with small angle x-ray diffraction approaches. Moreover, the combination of AML and AT with other drugs had no inhibitory effect on cholesterol crystal formation. Both the combination of AML with the HMG-CoA reductase inhibitor lovastatin and the combination of AT with the CCB nifedipine failed to interfere with cholesterol domain formation, as compared to control samples (Figure 4).

Cholesterol domains were very prominent in these samples with a unit cell periodicity of 34.0 Å. These discrete structures coexist with the surrounding sterol-poor region of the membrane. At 5°C and 74% relative humidity, the surrounding sterol-poor region of the membrane samples had the following d -space values: control (55.5 Å), AML/lovastatin (54.4 Å), and AT/nifedipine

(56.5 Å). Finally, when AML and AT were added separately to the cholesterol-enriched membrane samples, they did not interfere with domain formation.

The synergistic effect of AML and AT on cholesterol domain formation was also observed at a lower concentration of cholesterol. At a cholesterol to phospholipid mole ratio of 1.1:1, the drug combination effectively interfered with cholesterol crystallization within the membrane samples (Figure 5). By contrast, when used separately, the drugs had no effect on domain formation, even at this lower level of membrane cholesterol. At 5°C and 74% relative humidity, the surrounding sterol-poor region of the membrane samples had the following *d*-space values: control (55.5 Å), AML alone (54.4 Å), AT alone (55.8 Å), and AML/AT (53.9 Å).

An explanation for the synergistic effect of AML and AT on the organization of cholesterol may be their chemical properties. AML has very high lipophilicity as compared to other CCBs and a formal positive charge at physiologic pH. An electrostatic interaction between AML and AT as well as the phospholipid headgroup region of the membrane contributes to the high affinity of this agent for the lipid bilayer. Moreover, the charged amino-ethoxy function of AML directs the drug to a region of the membrane that overlaps the steroid nucleus of cholesterol molecules, an effect that may directly lead to a disruption in the self-association of cholesterol molecules in the membrane. Likewise, it has been observed that AT partitions to a similar location in the membrane as AML.

The key finding was the observation that the combination of AML and AT inhibited the formation of separate cholesterol domains in atherosclerotic-like membranes in a synergistic

fashion. This biophysical effect of the drug combination was directly characterized with small angle x-ray diffraction approaches using lipid membranes enriched with cholesterol. As cholesterol aggregates within the membrane may serve as nucleating sites for extracellular free cholesterol crystal formation in the vessel wall, the ability of the AML/AT combination to block such sterol domain formation indicates a novel antiatherosclerotic mechanism of action. This observed effect appears to be distinct for these drugs as other combinations failed to reproduce this change in the aggregation properties of free cholesterol.

In atherosclerosis, the incidence of lesion rupture and thrombosis is affected by the lipid composition of the atherosclerotic plaque. The lipid component of atherosclerotic lesions consists primarily of cholesterol and phospholipid, with lesser amounts of fatty acid and triacylglycerol. Over time, cholesterol forms crystalline structures in the human atheroma, an event that contributes to overall lesion mass and plaque instability. Once crystallized, cholesterol within the lesion is essentially inert and cannot be effectively removed by lipoprotein acceptors in the plasma. By contrast, non-crystallized cholesterol associated with foam cell membranes or intracellular stores can be depleted by plasma HDL and pharmacological interventions, leading to lesion regression.

Recent reports indicate that the cellular membrane is a cellular site for free cholesterol accumulation, leading to discrete sterol-rich domains and eventually crystal. In macrophage foam cells, for example, a critical mass of cholesterol is achieved following lipoprotein (native or oxidized) uptake and/or phagocytosis of lipid released from neighboring necrotic foam cells. Ultimately, a nucleating event will occur at a critical concentration of cholesterol enrichment,

leading to cholesterol domain development within the membrane. By interfering with the formation of highly organized cholesterol aggregates within the membrane, the combination of AML and AT may significantly slow or even prevent subsequent crystal development in the vessel wall, and thereby block the progression of an otherwise irreversible step in atherosclerosis. Moreover, these agents may work synergistically with HDL and lipid-lowering therapy in reducing the accumulation of cholesterol crystals in the wall of the diseased artery by maintaining cholesterol in a non-crystalline or dynamic state in cellular membranes.

The mechanism by which AML and AT interfere with the aggregation of cholesterol into discrete domains may be related to its their molecular membrane interactions. At physiologic pH, more than 90% of the amino ethoxy function associated with the #2 position of the dihydropyridine ring of AML is in the charged state. This positive charge contributes to specific electrostatic interactions of AML with phosphate groups associated with the phospholipid bilayer surface. The results of previous small-angle x-ray diffraction, differential scanning calorimetry and nuclear magnetic resonance analyses support a molecular model that places the charged amino function of AML near oppositely charged groups in the phospholipid headgroup region. Simultaneously, the hydrophobic portion of the dihydropyridine molecule is buried within the membrane hydrocarbon core, adjacent to the headgroup region. These biophysical measurements indicate that the time-averaged location of the ring structure for AML overlaps the sterol nucleus of cholesterol in the membrane, where it can then modulate certain biophysical effects of the molecule, and interfere with its self-association. Likewise, small-angle x-ray diffraction approaches demonstrated that AT partitioned to a discrete location in the membrane bilayer.

Thus, this unexpected, synergistic effect can be attributed to the molecular interactions of these compounds with membrane lipid constituents. This finding has important relevance for the treatment of coronary artery disease (CAD) as this disorder is characterized by the abnormal accumulation of free cholesterol into separate, membrane domains (d -space of 34.0 Å). These domains disrupt cellular function and lead to extracellular crystal formation, an important feature of the unstable atherosclerotic plaque. Small angle x-ray diffraction analyses demonstrated, for the first time, that the combination of AML and AT blocked the aggregation of free cholesterol into crystalline-like domains at low, nanomolar concentrations. By contrast, the combination of these agents with other related drugs showed no inhibitory effect on cholesterol crystal formation. These findings indicate that the combination of AML and AT produces a novel antiatherosclerotic effect by disrupting cholesterol crystal formation in atherosclerotic-like membranes. By disrupting the formation of cholesterol crystals in the vessel wall, the AML/AT combination would reduce plaque instability while facilitating cholesterol efflux to sterol acceptor particles, such as HDL. This new antiatherosclerotic mechanism of action for the AML/AT combination would complement the separate activities of these agents in the effective treatment of cardiovascular disease.

NO Release from Aortic Endothelial Cells: An amperogram (concentration of NO recorded versus time) obtained during amlodipine (5 µmol/L)- stimulated NO release from the single rabbit aorta endothelial cell is shown in Figure 6. NO was released in the form of sinusoidal pulse with maximum concentration of 120 ± 15 nmol/L. After the injection of atorvastatin, a small increase of NO concentration was also observed with maximum of 20 ± 5

nmol/L. This concentration is very low and may be due to the process of scavenging of superoxide by atorvastatin (i.e., an increase of net basal NO production). The injection of 5 μ mol/L of atorvastatin followed by the injection of 5 μ mol/L of amlodipine caused a massive release of NO reaching a maximum of 300 ± 10 mmol/L. The concentration of NO produced in association with the drug combination is twice what is expected based on the additive effects of the compounds, as recorded with separately with amlodipine or atorvastatin.

Figure 7 shows dose response curves for NO release stimulated by amlodipine, atorvastatin, and the mixture of 5 μ mol/L of amlodipine and variable concentrations (from 1 - 5 μ mol/L) of atorvastatin. Based on the data from this Figure, there is a significant synergistic effect observed after stimulation of NO release from endothelial cells by the combination of amlodipine and atorvastatin over a range of doses.

Therefore, the results of these analyses demonstrated a powerful synergistic effect for the combination of amlodipine and atorvastatin on the inhibition of cholesterol crystal formation and nitric oxide release from rabbit aortic endothelial cells. The results of this study provide compelling scientific support for the combined use of AML and AT in the treatment of cardiovascular disorders. These novel antiatherosclerotic effects of the AML/AT combination complement the separate activities of these agents in the treatment of cardiovascular disease, including CAD.

It will now be apparent to those skilled in the art that other embodiments, improvements, details, and uses can be made that are consistent with the letter and spirit of the foregoing disclosure and within the scope of this patent and the appended claims.

100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000